

24. Schwann TA, Habib RH, Zacharias A, Parenteau GL, Riordan CJ, Durham SJ, Engoren M. Effects of body size on operative, intermediate, and long-term outcomes after coronary artery bypass operation. *Ann Thorac Surg* 2001;71:521-30.
25. Reeves BC, Ascione R, Chamberlain MH, Angelini GD. Effect of body mass index on early outcomes in patients undergoing coronary artery bypass surgery. *J Am Coll Cardiol* 2003;42:668-76.
26. Potapov EV, Loebe M, Anker S, Stein J, Bondy S, Nasser BA, et al. Impact of body mass index on outcome in patients after coronary artery bypass grafting with and without valve surgery. *Eur Heart J* 2003;24:1933-41.
27. Uva MS, Rodrigues V, Monteiro N, Manuel Pedro A, Caria R, Mesquita A, et al. Cardiac surgery and morbid obesity. *Rev Port Cardiol* 2002;21:255-64.
28. Rockx MA, Fox SA, Stitt LW, Lehnhardt KR, McKenzie FN, Quantz MA, et al. Is obesity a predictor of mortality, morbidity and readmission after cardiac surgery? *Can J Surg* 2004;47:34-8.
29. Birkmeyer NJ, Charlesworth DC, Hernandez F, Leavitt BJ, Marrin CA, Morton JR, et al. Obesity and risk of adverse outcomes associated with coronary artery bypass surgery. Northern New England Cardiovascular Disease Study Group. *Circulation* 1998;97:1689-94.
30. Kuduvalli M, Grayson AD, Oo AY, Fabri BM, Rashid A. Risk of morbidity and in-hospital mortality in obese patients undergoing coronary artery bypass surgery. *Eur J Cardiothorac Surg* 2002;22:787-93.
31. Ranucci M, Cazzaniga A, Soro G, Morricone L, Enrini R, Caviezel F. Obesity and coronary artery surgery. *J Cardiothorac Vasc Anesth* 1999;13:280-4.
32. Fasol R, Schindler M, Schumacher B, Schlaudraff K, Hannes W, Seitelberger R, Schlosser V. The influence of obesity on perioperative morbidity: retrospective study of 502 aortocoronary bypass operations. *Thorac Cardiovasc Surg* 1992;40:126-9.
33. Moulton MJ, Creswell LL, Mackey ME, Cox JL, Rosenbloom M. Obesity is not a risk factor for significant adverse outcomes after cardiac surgery. *Circulation* 1996;94(Suppl):II87-92.
34. Kim J, Hammar N, Jakobsson K, Luepker RV, McGovern PG, Ivert T. Obesity and the risk of early and late mortality after coronary artery bypass graft surgery. *Am Heart J* 2003;146:555-60.
35. Brandt M, Harder K, Walluscheck KP, Schöttler J, Rahimi A, Möller F, Cremer J. Severe obesity does not adversely affect perioperative mortality and morbidity in coronary artery bypass surgery. *Eur J Cardiothorac Surg* 2001;19:662-6.
36. Jin R, Grunkemeier GL, Furnary AP, Handy JR, Jr. Is obesity a risk factor for mortality in coronary artery bypass surgery? *Circulation* 2005;111:3359-65.
37. Romero-Corral A, Montori VM, Somers VK, Korinek J, Thomas RJ, Allison TG, et al. Association of bodyweight with total mortality and with cardiovascular events in coronary artery disease: a systematic review of cohort studies. *Lancet* 2006;368:666-78.
38. Odrowaz-Sypniewska G. Markers of pro-inflammatory and pro-thrombotic state in the diagnosis of metabolic syndrome. *Adv Med Sci* 2007;52:246-50.
39. Hopkins TA, Ouchi N, Shiata R, Walsh K. Adiponectin actions in the cardiovascular system. *Cardiovasc Res* 2007;74:11-8.
40. Steppan CM, Bailey ST, Bhat S, Brown EJ, Banerjee RR, Wright CM, et al. The hormone resistin links obesity to diabetes. *Nature* 2001;409:307-12.
41. Steppan CM, Lazar MA. The current biology of resistin. *J Intern Med* 2004;255:439-47.
42. Mu H, Ohashi R, Yan S, Chai H, Yang H, Lin P, et al. Adipokine resistin promotes in vitro angiogenesis of human endothelial cells. *Cardiovasc Res* 2006;70:146-57.
43. Kougias P, Chai H, Lin PH, Lumsden AB, Yao Q, Chen C. Adipocyte-derived cytokine resistin causes endothelial dysfunction of porcine coronary arteries. *J Vasc Surg* 2005;41:691-8.

Submitted Jun 6, 2008; accepted Aug 8, 2008.

*Additional material for this article may be found online at [www.jvascsurg.org](http://www.jvascsurg.org).*

## DISCUSSION

**Dr Angeliki Vouyouka (New York, NY):** The paper was very interesting, but I have a few questions.

From the data that you show us here, the obese class I and the overweight patients compared to the normal and the underweight, they appeared to be healthier. They were more likely to be diabetic and hypertensive; however, they were less likely to be smokers, they have better ASA risk class, and they were less likely to have COPD. So do you think that part of your observation reflects some selection bias among the surgeons, since they were all elective cases, and therefore when they had to operate on obese patients they chose the ones that they were in best health as compared to the rest of them?

Secondly, you have insinuated the answer of this question, but for better clarification, what percentage of the obese patients had endovascular procedures, as compared to normal weight and underweight?

And thirdly, again, in the same spirit with the first question, in this data the obese patients, obese class I, and the overweight, they did have higher albumin levels, a fact that again perhaps reflects some selection bias. Do we know patients with low preoperative serum albumin, obese or overweight, how they did compare to the ones with high levels of serum albumin?

I want again to thank the committee for the privilege to discuss this very interesting paper.

**Dr Eleftherios Xenos:** To answer the second question, we don't have a breakdown between endovascular versus open repair of aneurysms. There is a procedure category that was labeled other and comprised approximately 30% of the patients. It included the peripheral balloon angioplasties and stenting procedures, but I

don't have an exact number of how many people had an endovascular versus open general vascular procedure.

And I agree with the observation that there might very well be a selection bias where relatively healthy obese subjects were offered an operation and very sick morbidly obese patients were denied an operation and that might be partially responsible for what we see.

**Dr Martin Veller (Johannesburg, South Africa):** Many of the individuals we treat with a low BMI have such a low BMI as a result of an underlying chronic disease. Were you in any way able to look at this in your study? For example, were the serum albumin levels helpful in differentiating those individuals with a low BMI as a result of a chronic disease from those with a low BMI without a chronic disease?

**Dr Xenos:** Yes, indeed, it does seem that if somebody is underweight, he's underweight for a reason. Many of our risk factors were highest in underweight patients including disseminated cancer and recent weight loss. There was also a higher incidence of patients with COPD. There was, however, no single etiology for being underweight and having chronic disease.

**Dr Starros Kakkos (Detroit, MI):** Do you think that slightly overweight people are naturally selected to be more resistant to trauma or similar conditions, like surgery?

**Dr Xenos:** I think it is hard to tell. In our study population, moderate overweight and obese I type patients had the least amount of risk factors.

**Dr Kakkos:** Probably because the definition and grades of obesity are based on survival statistics.

**Dr Xenos:** Yes, that is true, the patients were stratified according to their BMI, not survival statistics.